




Case Study: Seasonality Monitoring in Sweden & Finland



Northern high latitudes have been under close scrutiny by environmental researchers over the past few years. Such areas are particularly sensitive to changing climate, but we know relatively little about how they will respond, particularly in terms of the carbon and water cycles. Over the past two seasons, including the International Polar Year (IPY) in 2007-8, members of ABACUS (Arctic Biosphere-Atmosphere Coupling Across Multiple Scales), a multi-disciplinary team of researchers from Durham, Edinburgh, Sheffield York, University College, London (UCL), Macaulay, Stirling and CEH Wallingford, have been carrying out research at Abisko in Sweden, and Kevo in Finland.



The ABACUS project is exploring the links between the carbon and water cycles, and what the controls on these processes are at a range of scales. This work requires a wide range of measurements and models describing things such as vegetation amount and type, CO₂ and CH₄ fluxes from plants and soil, and detailed physiological processes controlling plant growth. The picture below shows one of four ABACUS instrument sites, two each in Sweden and Finland, which have been in place since 2006-7. This picture was taken in late August, when the vegetation is shutting down photosynthetic activity for the long winter ahead.

The second picture shows the same site, with a large reindeer encroaching onto the experimental plot. The Abisko area is part of the migratory grazing route for the reindeer herds of the indigenous Sami people who live across the region.

The two graphs below show the results from Skye sensors mounted at two locations in Abisko in 2007 and 2008, one a mountainous tundra site (seen in figures 1 and 2), and one at a lower, forested site. The sensors measure incident and reflected radiation in visible red and near-infrared channels corresponding to those of the Landsat satellite. From these measurements we can calculate Normalised Difference Vegetation Index (NDVI), which is related to the amount and greenness of vegetation. The graphs illustrate the different seasonal responses of the tundra and forest vegetation and the rapid green-up of the tundra following snow melt in late May (where NDVI is close to zero), compared with the slower, later development of the forested site. There are also subtle changes in the timing of these events from 2007 to 2008.

These measurements are being combined with airborne and satellite data and other ground-based measurements to give a broader picture of the biophysical response of these landscapes through the seasons.

ABACUS is funded by the Natural Environment Research Council (NERC). More information on the ABACUS project can be found at www.abacus-ipy.org.



Acknowledgements and Contacts

We would like to thank Dr. M. Disney & Dr. A. Prieto-Blanco of UCL Department of Geography for supplying us with a case study.

Skye Instruments Ltd

21, Ddole Enterprise Park, Llandrindod Wells, Powys LD1 6DF, United Kingdom
TEL: +44 (0)1597 824811 EMAIL: skyeemail@skyeinstruments.com WEB: www.skyeinstruments.com